بسم الله الرحمن الرحيم

Cell Injury

By

Dr. Dina Abdallah


Glycocalyx
Microvilli
Phagosomes
Lysosome
Polysome
Rough endoplasmic reticulum(RER)
Smooth endoplasmic reticulum(SER)
Free ribosome
Glycogen
Cilia
Zonula occludens
Zonula adherens
Microfilaments
Golgi complex
Centriole
Nucleus
Nucleolus
Mitochondrion
Basement membrane
Collagen

Structure of cells & related functions

I- Nucleus:-

Replication of DNA, production of mRNA and synthesis of nuclear proteins.

It also contains :-

Nucleolus represent new synthesized ribosomes + rRNA → direct synthesis of specific proteins in cytoplasm.

II- Cytoplasmic organelles:-

***Mitochondria :** major source of ATP→ energy

***E R** rough → protein

Smooth → drug intoxication

***Golgi apparatus :** in secreting cells (package of selective protein from ER)

***Lysosomes** (hydrolytic enzymes **e.g.** acid hydrolyses)

III- Cell membrane

- Maintain integrity of cells .
- Active Na pump.
- Intake of exogenous material either phagocytosis or pinocytosis
- Receptors.


Smooth Endoplasmic Reticulum
Nucleus
Microfilaments
Golgi Complex
Mitochondrion
Lysosome
Nucleolus
Chromatin
Rough Endoplasmic Reticulum
Microtubules
Vesicle
Plasma Membrane

Homeostasis

Normal cells have a fairly narrow range of function or steady state

Injurious agents

Adaptation

-Hypertrophy
-Atrophy
-Hyperplasia
-Metaplasia

Cell injury

Reversible

-Cloudy swelling
-Hydropic degeneration
-Fatty changes

Irreversible

Necrosis

Apoptosis

Intra and Extracellular Accumulations and Depositions

- Mucin
- Gout
- Hyalinosis
- Pathological pigmentation
- Amyloidosis
- Pathological calcification

Exposure of the cell to excess physiologic or pathologic stress leads to:

A. Adaptation.

- It is a modification of cell morphology and function to achieve a new steady but altered state preserving the viability of cells.
- *a. Hypertrophy* (increase in the size of the cells e.g. skeletal muscle cells in body builders).
- *b. Atrophy* (decrease in the size of the cells e.g. atrophy of genital organs after menopause).
- *c. Hyperplasia* (increase in the number of cells e.g. hyperplasia of the breast in response to estrogenic stimulation).
- *d. Metaplasia* (change of the type of the cell to another type of the same category e.g. squamous metaplasia of urothelial cells in case of bilharziasis).

Hypertrophy





Individual cells
become larger

e.g. muscle hypertrophy after regular exercise

©Stevens&Lowe 1999

Normal breast

Lactating breast


Basement membrane
Normal columnar epithelium
Reserve cells
Squamous metaplasia
A

B. Cell injury:

- - It is a change in the cell morphology and function in response to stress (irritant). This occurs when the cells can not able to adapt.
- - The form of cell injury depends on:

a- Type, duration and severity of the irritant

b- Type of injured tissue (its adaptability power)

e.g. brain tissue is sensitive to hypoxia within2-5 minutes while cardiac ms can adapt for 2-6 hours

* Causes of cell injury:

- Oxygen deprivation: hypoxia, ischemia.
- Oxygen free radicals.
- Physical agents: heat, cold, irradiation.
- Chemical agents: drugs, toxins.
- Infection (bacterial, viral,…….).
- Immunologic reaction.
- Genetic causes.
- Nutritional imbalance e.g. starvation.

* These injurious agents can affect any structural component of the cell but the mitochondrial injury is the most accepted theory that can explain most forms of cell injury.

A. REVERSIBLE CELL INJURY

CLOUDY SWELLING

- ***Def:*** mild degeneration characterized by cell swelling and granularity of the cytoplasm due to intracellular accumulation of water.
- **** Etiology:*** caused by mild irritant acting for short duration.

*Morphological features:

**** Organs affected:*** parenchymatous organs e.g. liver, kidney & heart…

**** N/E:*** the affected organ shows:

- ***Size:*** enlarged (swollen).
- ***Shape:*** preserved.
- ***Surface:*** smooth.
- ***Color:*** pale.
- ***Consistency:*** soft.
- ***Capsule:*** stretched (non-adherent)***.***
- ***C/S:*** bulges outwards with rounded edges.

**** M/E:***

- Swollen granular cytoplasm with intact nucleus.
- The capillaries between the cells are compressed

VACUOLAR & HYDROPIC DEGENERATION

***Def.** Advanced stage of cloudy swelling characterized by excess water accumulation inside the cells in the form of cytoplasmic vacuoles (vacuolar) or as large space (hydropic).

***Etiology:** caused by mild injurious agent acting for more prolonged duration

*** Examples:**

- Liver in: alcoholism – viral hepatitis-CCL4.
- Skin: burn, urticaria.
- Pancreas: DM

******Morphological features:*** as cloudy swelling.



Cellular swelling (hydropic change), liver

FATTY CHANGE (Steatosis)

***Def.** . Intracellular accumulation of excess neutral fat (triglycerides) in parenchymatous organs.

***Etiology:** caused by mild injurious agent acting for more prolonged duration

*Morphological features:

- ***Organs affected:*** liver, heart.

**** N/E:*** the affected organ shows:

- ***Size:*** enlarged .
- ***Shape:*** preserved.
- ***Surface:*** smooth.
- ***Color:*** yellowish.
- ***Consistency:*** soft & greasy.
- ***Capsule:*** stretched (non-adherent).
- ***C/S:*** bulges outwards with rounded edges.

**** M/E:***

- The cells are swollen, the cytoplasm contains droplets of fat (in mild cases) termed microvesicular steatosis, or vacuoles pushing the nucleus to one side in severe cases (signet ring appearance).


Fatty liver

A



B



C

Staining properties:

- Formalin fixed and usual processing → unstained vacuoles.

- Fresh and frozen sections and stained with:

Osmic acid

Sudan III

Sudan black






Sudan III



B. IRREVERSIBLE CELL INJURY

Necrosis

Def: local death of a group of cells within the living body.

Etiology: *severe irritant.*

**** N/E:*** The necrotic area is:

- Well-defined.
- Swollen.
- Opaque.
- Pale yellow or pale red.
- Surrounded by a red zone of inflammatory hyperemia.



coagulative necrosis (infarction) in the renal cortex of the kidney

* Types of necrosis:

1. Coagulative necrosis. E.g. ischaemia of the heart (MI)
2. Liquefactive necrosis. E.g. abscess
3. Caseation necrosis. TB
4. Fat necrosis include traumatic (breast) and enzymatic (pancratitis)
5. Zenker's necrosis. E.g. typhyoid fever
6. Fibrinoid necrosis. E.g. rheumatic fever








CENTIMETERS



APOPTOSIS

**** Def:*** programmed single cell death.

****Types:***

- I. Physiological apoptosis.
- During embryogenesis.
- During menstruation.
- Menopause.
- II. Pathological apoptosis.
- Viral hepatitis.
- Neoplasia.
- Graft rejection.

Differences between apoptosis & necrosis

Necrosis	Apoptosis
•Groups of cells or tissue •No gene activation. •Mediated by calcium-activated lysosomal enzymes. •Decreased ATP •Swelling of cells . •Cellular membranes ruptured early. •Inflammatory reaction around •Always pathological	•Single cell. •Genetic activation •Mediated by genetically controlled cytoplasmic enzymes •Normal ATP •Shrunken cells •Cellular membranes intact until late when apoptotic bodies separate. •No inflammation. •Could be physiological or pathological